


## Answers Tutorials 2 Thermodynamics 2, 2025/2026

### Exercise 5

The corrected phase diagram is shown in the figure below, with the corrections in red. The 5 mistakes are:

1.  $P$  and  $T$  have been swapped
2.  $l$  and  $g$  have been swapped
3. 'sublimation point' should be 'triple point'
4. the critical point is where the  $l, g$  phase boundary line ends
5. the error in the red circle needs clarification. In the red circle four phase boundary lines come together in a single point. This is not possible according to Gibbs' phase rule,  $F = C - P + 2$ , as four phases ( $s_3, s_4, s_5$  and  $l$  would be mutually in equilibrium, so  $P = 4$ , while  $C = 1$  because we are dealing with a pure compound (with a unary phase diagram). This would result in  $F = C - P + 2 = 1 - 4 + 2 = -1$ , so not allowed; A pure compound cannot have four phases in mutual equilibrium.



### Exercise 6

We assume that the vapour behaves as a perfect gas, and that the enthalpy of vaporization is independent of the temperature in the pressure interval. With these assumptions the Clausius-Clapeyron equation becomes:

$$P = P^* \exp(-\chi) \quad \text{with} \quad \chi = \frac{\Delta_{\text{vap}}H}{R} \left( \frac{1}{T} - \frac{1}{T^*} \right) \quad \text{which implies}$$

$$\ln \frac{P^*}{P} = \chi,$$

or

$$\frac{1}{T} = \frac{1}{T^*} + \frac{R}{\Delta_{\text{vap}}H} \ln \frac{P^*}{P} = \frac{1}{297.25} + \left( \frac{8.314}{28.7 \cdot 10^3} \right) \ln \frac{400}{500} = 3.30 \cdot 10^{-3} \text{ K}^{-1},$$

so  $T = 303 \text{ K} = 30 \text{ }^\circ\text{C}$ .

## Exercise 7

The three phase boundary lines intersect in the triple point. These lines can be found using the following equations between  $P$  and  $T$  for the phase boundaries in a  $(P, T)$ -diagram. ( $(P, T)$  and  $(P^*, T^*)$  correspond to two points on a phase boundary line):

$$P = P^* + \frac{\Delta_{\text{fus}}H}{\Delta_{\text{fus}}V} \ln \frac{T}{T^*} \quad (\text{s} - \text{l}),$$

$$P = P^* \exp(-\chi); \quad \chi = \frac{\Delta_{\text{vap}}H}{R} \left( \frac{1}{T} - \frac{1}{T^*} \right) \quad (\text{l} - \text{g}) \text{ and}$$

$$P = P^* \exp(-\chi); \quad \chi = \frac{\Delta_{\text{sub}}H}{R} \left( \frac{1}{T} - \frac{1}{T^*} \right) \quad (\text{s} - \text{g}),$$

for which we used the assumption that  $\Delta H$  and  $\Delta V$  hardly change in the temperature range around the triple point.

With this assumption we can calculate  $\Delta_{\text{sub}}H$  using  $\Delta_{\text{sub}}H = \Delta_{\text{fus}}H + \Delta_{\text{vap}}H = 10.6 + 30.8 = 41.4 \text{ kJ/mol}$ .

$\Delta_{\text{fus}}V$  follows from (benzene  $\text{C}_6\text{H}_6$ ,  $M(\text{C}_6\text{H}_6) = 78.11 \text{ g/mol}$ )

$$\Delta_{\text{fus}}V = \frac{M}{\rho(\text{l})} - \frac{M}{\rho(\text{s})} = \frac{78.11}{0.879} - \frac{78.11}{0.891} = 1.197 \text{ cm}^3/\text{mol}.$$


Entering these values and changing all of them to SI (mksA) units, except for the pressures, which we leave in Torr (1 Torr = 133.322 Pa), we can find the three phase boundary lines by plugging in the temperatures in Kelvin. For  $(T^*, P^*)$  we can choose the triple point values (278.65 K, 36 Torr).

$$P = P^* + \frac{10.6 \cdot 10^3}{1.197 \cdot 10^{-6}} \ln \frac{T}{T^*} = P^* + 8.855 \cdot 10^9 [\text{Pa}] \ln \frac{T}{T^*} = P^* + 6.64 \cdot 10^7 [\text{Torr}] \ln \frac{T}{T^*} \quad (\text{s} - \text{l}),$$

$$P = P^* \exp(-\chi); \quad \chi = \frac{30.8 \cdot 10^3}{8.314} \left( \frac{1}{T} - \frac{1}{T^*} \right) = 3705 [\text{K}] \left( \frac{1}{T} - \frac{1}{T^*} \right) \quad (\text{l} - \text{g}) \text{ en}$$

$$P = P^* \exp(-\chi); \quad \chi = \frac{41.4 \cdot 10^3}{8.314} \left( \frac{1}{T} - \frac{1}{T^*} \right) = 4980 [\text{K}] \left( \frac{1}{T} - \frac{1}{T^*} \right) \quad (\text{s} - \text{g}).$$

The three lines can be found in the figure below as  $a$ ,  $b$  en  $c$ , around the triple point of  $P^* = 36 \text{ Torr}$  (4.800 kPa) and  $T^* = 5.50 \text{ }^\circ\text{C}$  (278.65 K). The boundary line of  $c$  for  $T > T^*$  (dashed line) is metastable, and the same applies to  $b$  for  $T < T^*$ .



## Exercise 8

The change in freezing point as a result of the higher pressure in the mercury column can be calculated with the Clapeyron equation (use  $\Delta_{\text{fus}}G = \Delta_{\text{fus}}H - T_{\text{fus}}\Delta_{\text{fus}}S = 0$  for a phase transition (so at  $T = T_{\text{fus}}$ )).

$$\frac{dP}{dT} = \frac{\Delta_{\text{fus}}S}{\Delta_{\text{fus}}V} = \frac{\Delta_{\text{fus}}H}{T_{\text{fus}}\Delta_{\text{fus}}V}, \quad \text{so} \quad dP = \frac{\Delta_{\text{fus}}H}{T_{\text{fus}}\Delta_{\text{fus}}V}dT \quad \text{and} \quad \int_{P_1}^{P_2} dP = \int_{T_{\text{fus}}}^{T_{\text{fus},2}} \frac{\Delta_{\text{fus}}H}{T_{\text{fus}}\Delta_{\text{fus}}V}dT.$$

The Clapeyron equation describes the pressure change as a function of the temperature as long as we stay on the phase boundary line, which means that we should actually write  $dP_{\text{fus}}$  and  $dT_{\text{fus}}$  instead of  $dP$  and  $dT$ . We assume that for the small temperature change all quantities in the integral (except for  $T_{\text{fus}}$  which changes due to the pressure in the mercury column) are constant, which leads to

$$P_2 - P_1 = \frac{\Delta_{\text{fus}}H}{\Delta_{\text{fus}}V} \int_{T_{\text{fus}}}^{T_{\text{fus},2}} \frac{dT}{T} = \frac{\Delta_{\text{fus}}H}{\Delta_{\text{fus}}V} \ln \frac{T_{\text{fus},2}}{T_{\text{fus}}}.$$

The total pressure difference over the column is  $\Delta P = (P_2 - P_1) = \rho(l)g\Delta h$ , with  $\Delta h$  the column height and  $g = 9.81 \text{ m/s}^2$ , which results in a change in freezing point at the bottom of the column determined by

$$\ln \frac{T_{\text{fus},2}}{234.3} = (P_2 - P_1) \frac{\Delta_{\text{fus}}V}{\Delta_{\text{fus}}H} = \rho(l)g\Delta h \frac{\Delta_{\text{fus}}V}{\Delta_{\text{fus}}H} = 13.6 \cdot 10^3 \cdot 9.81 \cdot 10.0 \frac{0.517 \cdot 10^{-6}}{2.292 \cdot 10^3} = 3.0094 \cdot 10^{-4}.$$

This results in a temperature  $T_{\text{fus},2} = 234.4 \text{ K}$ , so an increase of  $0.1 \text{ K}$ .

The answer therefore is that the bottom of the column just freezes at a temperature of  $234.4 \text{ K}$ .

## Exercise 9

- a) The suggested Maxwell relation with a term  $\left(\frac{\partial V}{\partial T}\right)_P = V\alpha$  follows from  $dG = VdP - SdT$ .

$G$  is a state function with independent variables  $P$  and  $T$ , which means that

$$\frac{\partial^2 G}{\partial T \partial P} = \frac{\partial^2 G}{\partial P \partial T}, \quad \text{and therefore} \quad \left(\frac{\partial V}{\partial T}\right)_P = -\left(\frac{\partial S}{\partial P}\right)_T.$$

According to this relation  $\left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial V}{\partial T}\right)_P = -V\frac{1}{V}\left(\frac{\partial V}{\partial T}\right)_P = -V\alpha$ , and therefore  $dS|_T = -V\alpha dP|_T$ .

The volume is  $V = \frac{M}{\rho}n = \frac{200}{2 \cdot 10^3} \cdot 1 = 0.1 \text{ l}$  for  $1 \text{ mol}$ .

If we assume that  $\alpha$  is independent of the pressure we find (note the units)

$$\Delta S = -V\alpha\Delta P = -0.1 \cdot 2 \cdot 10^{-3}(100 - 1) = -0.02 \text{ l} \cdot \text{bar} \cdot \text{K}^{-1} = -0.02 \cdot 10^2 \text{ m}^3 \text{Pa} \cdot \text{K}^{-1} = -2 \text{ J} \cdot \text{K}^{-1}$$

- b)  $\Delta U = W + Q = -\int P_{\text{ext.}}dV + Q \approx Q$  because the volume change is negligible.  
In that case the difference between a reversible and an irreversible process is also negligible, and since  $U$  is a state function we find:  
 $\Delta U = Q^{\text{rev}} = T\Delta S = -300 \cdot 2 = -0.6 \text{ kJ}$ .
- c)  $dH = dU + PdV + VdP = dQ + VdP$ , so  
 $\Delta H = Q + \int VdP \approx Q + V\Delta P$  because the volume change is negligible, so  
 $\Delta H = -600 + 0.1 \cdot 10^{-3}(100 - 1) \cdot 10^5 \text{ J} = 0.4 \text{ kJ}$ .
- d)  $dA = dU - TdS - SdT = -PdV - SdT$ , so  
 $\Delta A \approx 0$  because it is an isothermal process and the volume change is negligible.
- e)  $dG = dA + PdV + VdP = -SdT + VdP$ , so  
 $\Delta G \approx V\Delta P$  because it is an isothermal process and the volume change is negligible.  
 $\Delta G = 0.1 \cdot 10^{-3}(100 - 1) \cdot 10^5 \text{ J} = 1 \text{ kJ}$ .